

## Student Session Topic: Related Rate Problems

Related Rate problems appear only occasionally on the AP calculus exams. Typically there will be a straightforward question in the multiple-choice section; on the free-response section a related rate question will be part of a longer question or.

The thing with related rate questions is that there are, apparently, only a limited number of situations in which they appear – cones or cylinders being filled or emptied, people walking away from lampposts, ladders sliding down walls, vehicles moving in perpendicular directions, etc. These appear in all the textbooks and it is difficult to find suitable new questions.

### What you should be able to do:

- Read, understand and translate the verbal description into symbols relating the variables that are changing.
- Understand what quantities are changing and which are constant throughout the problem.
- Understand the geometry of the situation. Use geometric relationships to replace a variable with another. This may require working with similar figures.
- Do implicit differentiation with respect to time.
- A common misunderstanding is to substitute the specific values of variables into the equation before differentiating.
  - Any quantity that does not change during the course of the problem can be substituted at the beginning
  - Any quantity that changes should not be substituted until the derivative is being evaluated at the specific time stated in the question.

### Questions:

Free-response questions for the prep session: 2002 AB 5 (no calculator), 2008 AB 3 and 2003 AB 5/BC 5.

**WATCH and LISTEN** to the multiple-choice questions being solved

Go to <http://tinyurl.com/NMSI-Math-3> Click on the "Full Screen" arrow.  
Then click anywhere on the page to see and hear from that point on.

Click anywhere to go back anytime.

Answers: 1 A, 2 E, 3 B, 4 D, 5 C.


**Related Rates****Multiple Choice**

Identify the choice that best completes the statement or answers the question.

1. A railroad track and a road cross at right angles. An observer stands on the road 70 meters south of the crossing and watches an eastbound train traveling at 60 meters per second. At how many meters per second is the train moving away from the observer 4 seconds after it passes through the intersection?

a. 57.60  
b. 57.88  
c. 59.20  
d. 60.00  
e. 67.40

2.



In the triangle shown above, if  $\theta$  increases at a constant rate of 3 radians per minute, at what rate is  $x$  increasing in units per minute when  $x$  equals 3 units?


a. 3  
b.  $\frac{15}{4}$   
c. 4  
d. 9  
e. 12

3. The radius of a circle is decreasing at a constant rate of 0.1 centimeter per second. In terms of the circumference  $C$ , what is the rate of change of the area of the circle, in square centimeters per second?

a.  $-(0.2)\pi C$   
b.  $-(0.1)C$   
c.  $-\frac{(0.1)C}{2\pi}$   
d.  $(0.1)^2 C$   
e.  $(0.1)^2 \pi C$

4. If the base  $b$  of a triangle is increasing at a rate of 3 inches per minute while its height  $h$  is decreasing at a rate of 3 inches per minute, which of the following must be true about the area  $A$  of the triangle?
- a.  $A$  is always increasing.
  - b.  $A$  is always decreasing.
  - c.  $A$  is decreasing only when  $b < h$ .
  - d.  $A$  is decreasing only when  $b > h$ .
  - e.  $A$  remains constant.
5. The radius of a circle is increasing at a constant rate of 0.2 meters per second. What is the rate of increase in the area of the circle at the instant when the circumference of the circle is  $20\pi$  meters?
- a.  $0.04\pi \text{ m}^2/\text{sec}$
  - b.  $0.4\pi \text{ m}^2/\text{sec}$
  - c.  $4\pi \text{ m}^2/\text{sec}$
  - d.  $20\pi \text{ m}^2/\text{sec}$
  - e.  $100\pi \text{ m}^2/\text{sec}$

2002 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS



5. A container has the shape of an open right circular cone, as shown in the figure above. The height of the container is 10 cm and the diameter of the opening is 10 cm. Water in the container is evaporating so that its depth  $h$  is changing at the constant rate of  $-\frac{3}{10}$  cm/hr.


(Note: The volume of a cone of height  $h$  and radius  $r$  is given by  $V = \frac{1}{3}\pi r^2 h$ .)

- (a) Find the volume  $V$  of water in the container when  $h = 5$  cm. Indicate units of measure.
  - (b) Find the rate of change of the volume of water in the container, with respect to time, when  $h = 5$  cm. Indicate units of measure.
  - (c) Show that the rate of change of the volume of water in the container due to evaporation is directly proportional to the exposed surface area of the water. What is the constant of proportionality?
-

# AP<sup>®</sup> CALCULUS AB 2002 SCORING GUIDELINES

## Question 5

A container has the shape of an open right circular cone, as shown in the figure above. The height of the container is 10 cm and the diameter of the opening is 10 cm. Water in the container is evaporating so that its depth  $h$  is changing at the constant rate of  $-\frac{3}{10}$  cm/hr.



(The volume of a cone of height  $h$  and radius  $r$  is given by  $V = \frac{1}{3}\pi r^2 h$ .)

- Find the volume  $V$  of water in the container when  $h = 5$  cm. Indicate units of measure.
- Find the rate of change of the volume of water in the container, with respect to time, when  $h = 5$  cm. Indicate units of measure.
- Show that the rate of change of the volume of water in the container due to evaporation is directly proportional to the exposed surface area of the water. What is the constant of proportionality?

(a) When  $h = 5$ ,  $r = \frac{5}{2}$ ;  $V(5) = \frac{1}{3}\pi\left(\frac{5}{2}\right)^2 5 = \frac{125}{12}\pi \text{ cm}^3$

(b)  $\frac{r}{h} = \frac{5}{10}$ , so  $r = \frac{1}{2}h$   
 $V = \frac{1}{3}\pi\left(\frac{1}{4}h^2\right)h = \frac{1}{12}\pi h^3$ ;  $\frac{dV}{dt} = \frac{1}{4}\pi h^2 \frac{dh}{dt}$   
 $\left.\frac{dV}{dt}\right|_{h=5} = \frac{1}{4}\pi(25)\left(-\frac{3}{10}\right) = -\frac{15}{8}\pi \text{ cm}^3/\text{hr}$

OR

$$\frac{dV}{dt} = \frac{1}{3}\pi\left(r^2 \frac{dh}{dt} + 2rh \frac{dr}{dt}\right); \frac{dr}{dt} = \frac{1}{2} \frac{dh}{dt}$$

$$\left.\frac{dV}{dt}\right|_{h=5, r=\frac{5}{2}} = \frac{1}{3}\pi\left(\left(\frac{25}{4}\right)\left(-\frac{3}{10}\right) + 2\left(\frac{5}{2}\right)5\left(-\frac{3}{20}\right)\right)$$

$$= -\frac{15}{8}\pi \text{ cm}^3/\text{hr}$$

(c)  $\frac{dV}{dt} = \frac{1}{4}\pi h^2 \frac{dh}{dt} = -\frac{3}{40}\pi h^2$   
 $= -\frac{3}{40}\pi(2r)^2 = -\frac{3}{10}\pi r^2 = -\frac{3}{10} \cdot \text{area}$

The constant of proportionality is  $-\frac{3}{10}$ .

1 :  $V$  when  $h = 5$

5 :  $\left\{ \begin{array}{l} 1 : r = \frac{1}{2}h \text{ in (a) or (b)} \\ \\ V \text{ as a function of one variable} \\ \text{in (a) or (b)} \\ \\ \text{OR} \\ \frac{dr}{dt} \end{array} \right.$

2 :  $\frac{dV}{dt}$   
 $< -2 >$  chain rule or product rule error

1 : evaluation at  $h = 5$

2 :  $\left\{ \begin{array}{l} 1 : \text{shows } \frac{dV}{dt} = k \cdot \text{area} \\ 1 : \text{identifies constant of proportionality} \end{array} \right.$

units of  $\text{cm}^3$  in (a) and  $\text{cm}^3/\text{hr}$  in (b)

1 : correct units in (a) and (b)

Oil is leaking from a pipeline on the surface of a lake and forms an oil slick whose volume increases at a constant rate of 2000 cubic centimeters per minute. The oil slick takes the form of a right circular cylinder with both its radius and height changing with time. (Note: The volume  $V$  of a right circular cylinder with radius  $r$  and height  $h$  is given by  $V = \pi r^2 h$ .)

- (a) At the instant when the radius of the oil slick is 100 centimeters and the height is 0.5 centimeter, the radius is increasing at the rate of 2.5 centimeters per minute. At this instant, what is the rate of change of the height of the oil slick with respect to time, in centimeters per minute?
- (b) A recovery device arrives on the scene and begins removing oil. The rate at which oil is removed is  $R(t) = 400\sqrt{t}$  cubic centimeters per minute, where  $t$  is the time in minutes since the device began working. Oil continues to leak at the rate of 2000 cubic centimeters per minute. Find the time  $t$  when the oil slick reaches its maximum volume. Justify your answer.
- (c) By the time the recovery device began removing oil, 60,000 cubic centimeters of oil had already leaked. Write, but do not evaluate, an expression involving an integral that gives the volume of oil at the time found in part (b).

- (a) When  $r = 100$  cm and  $h = 0.5$  cm,  $\frac{dV}{dt} = 2000$  cm<sup>3</sup>/min  
and  $\frac{dr}{dt} = 2.5$  cm/min.

$$\frac{dV}{dt} = 2\pi r \frac{dr}{dt} h + \pi r^2 \frac{dh}{dt}$$

$$2000 = 2\pi(100)(2.5)(0.5) + \pi(100)^2 \frac{dh}{dt}$$

$$\frac{dh}{dt} = 0.038 \text{ or } 0.039 \text{ cm/min}$$

$$4 : \begin{cases} 1 : \frac{dV}{dt} = 2000 \text{ and } \frac{dr}{dt} = 2.5 \\ 2 : \text{expression for } \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$$

- (b)  $\frac{dV}{dt} = 2000 - R(t)$ , so  $\frac{dV}{dt} = 0$  when  $R(t) = 2000$ .

This occurs when  $t = 25$  minutes.

Since  $\frac{dV}{dt} > 0$  for  $0 < t < 25$  and  $\frac{dV}{dt} < 0$  for  $t > 25$ ,

the oil slick reaches its maximum volume 25 minutes after the device begins working.

$$3 : \begin{cases} 1 : R(t) = 2000 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$


- (c) The volume of oil, in cm<sup>3</sup>, in the slick at time  $t = 25$  minutes is given by  $60,000 + \int_0^{25} (2000 - R(t)) dt$ .

$$2 : \begin{cases} 1 : \text{limits and initial condition} \\ 1 : \text{integrand} \end{cases}$$

**AP® CALCULUS AB**  
**2003 SCORING GUIDELINES**

**Question 5**

A coffeepot has the shape of a cylinder with radius 5 inches, as shown in the figure above. Let  $h$  be the depth of the coffee in the pot, measured in inches, where  $h$  is a function of time  $t$ , measured in seconds. The volume  $V$  of coffee in the pot is changing at the rate of  $-5\pi\sqrt{h}$  cubic inches per second. (The volume  $V$  of a cylinder with radius  $r$  and height  $h$  is  $V = \pi r^2 h$ .)



- (a) Show that  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$ .
- (b) Given that  $h = 17$  at time  $t = 0$ , solve the differential equation  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$  for  $h$  as a function of  $t$ .
- (c) At what time  $t$  is the coffeepot empty?

(a)  $V = 25\pi h$

$$\frac{dV}{dt} = 25\pi \frac{dh}{dt} = -5\pi\sqrt{h}$$

$$\frac{dh}{dt} = \frac{-5\pi\sqrt{h}}{25\pi} = -\frac{\sqrt{h}}{5}$$

$$3 : \begin{cases} 1 : \frac{dV}{dt} = -5\pi\sqrt{h} \\ 1 : \text{computes } \frac{dV}{dt} \\ 1 : \text{shows result} \end{cases}$$

(b)  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$

$$\frac{1}{\sqrt{h}} dh = -\frac{1}{5} dt$$

$$2\sqrt{h} = -\frac{1}{5}t + C$$

$$2\sqrt{17} = 0 + C$$

$$h = \left(-\frac{1}{10}t + \sqrt{17}\right)^2$$

$$5 : \begin{cases} 1 : \text{separates variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition } h = 17 \\ \text{when } t = 0 \\ 1 : \text{solves for } h \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

(c)  $\left(-\frac{1}{10}t + \sqrt{17}\right)^2 = 0$

1 : answer

$$t = 10\sqrt{17}$$